

TOWARD THE NEW GRAVITATIONAL NONCOMMUTATIVE MECHANICS AND STATISTICAL MECHANICS OF QUANTUM BLACK HOLES

P. O. MAZUR

*Department of Physics and Astronomy
University of South Carolina
Columbia, SC 29208, USA*

This is a short account of our work on the statistical mechanics of ‘cold’ quantum black holes in the constituent model of a black hole^{3 4 6}. A quantum Schwarzschild black hole consists of *gravitational atoms* of Planckian mass scale^{3 4 6}. The gapless bosonic collective excitations of a bound state of N *gravitational atoms* dominate the thermodynamics of a cold quantum black hole. It turns out that it is only in the limit of large N , with the observable mean values of the gravitational mass-energy kept fixed, that we recover the Bekenstein thermodynamics¹ of spin-zero black holes. This is also the limit of the Boltzmann statistics.

1 On the Statistical Mechanics of Quantum Schwarzschild Black Holes

We have shown that a gravitating mass M in thermal equilibrium behaves statistically like a system of some number N of harmonic oscillators whose zero-point energy $\frac{\epsilon}{2}$ depends on N universally in such a way that $N\epsilon^2 \sim \frac{hc^5}{G}$, where G is the Newton constant, c is the velocity of light in vacuum, and h is the Planck constant. The large number N is also the number of weakly interacting gravitational atoms^{4 6 9} which are the constituents of a spin-zero black hole. The sum over all oscillators of the squares of zero-point energies is fixed and independent of the number of those oscillators. The enigmatic Bekenstein entropy of black holes was not yet derived on the basis of microscopic theory. Our work may be considered a first step in direction of presenting such a basis^{3 4 5 8 9}. The problem with all present approaches to this problem has been the silent assumption that the total entropy of gravitating systems (black holes) *must* be given by the Bekenstein formula¹ $S_{bh} = 4k\pi M^2$, where the mass M is in Planck units. The postulate of gravitational constituents (gravitational atoms) and gravitational oscillators (quanta) leads to Bekenstein formula only after a part of mass-energy fluctuations is neglected^{4 6}. The most unusual character of the gravitational mass-energy oscillators (quanta) is that they somehow manage, via the quadratic sum rule defining the Newton constant G , $\sum_i \epsilon_i^2 = b \frac{hc^5}{G}$, to reduce their zero-point energy when the number N of gravitational atoms grows⁴. The formula: $M^2 = \frac{\mu^2}{2\pi} N$, where $\mu^2 = \frac{hc}{G}$, was derived long time ago (1986) by the present author. This also means that a cold large gravitational mass $M \sim \mu\sqrt{N}$ consists of N *constituents*. The physical meaning of the ‘phenomenological’ entropy of Bekenstein¹ is that it is the measure of the number N of *constituents* making up a very ‘cold’ large body. The zero-point energy ϵ_i of *gravitational quanta* for a very large ‘cold’ mass is of an order of the Hawking thermal energy of quanta², $\epsilon_i \sim \frac{\mu^2}{(4\pi)^2 M}$. The more massive is a

gravitating mass the softer are the *gravitational quanta*. The number N of *constituent gravitational atoms* of a given spin-zero quantum Schwarzschild black hole determine the energy of quasi-thermal quanta and the Bekenstein-Hawking entropy, $kT_{bh} \sim \frac{\mu}{\sqrt{N}}$, $S_{bh} \sim kN$, but it is valid only in the particular limit when the interference terms are neglected⁴. Otherwise, as usual with oscillators, there are two sources of statistical fluctuations of mass-energy corresponding to the corpuscular and wave aspect of quanta⁴⁻⁶. We calculate the energy fluctuations of the system^a considered in⁴ in terms of the mean energy $\overline{E}$, where $\overline{E} = E_0 \cotanh(\frac{\epsilon}{2kT})$, $\epsilon = \frac{E_0}{N}$, $E_0^2 = \frac{b\mu^2 c^4 N}{4}$, and $\overline{(\Delta E)^2} = N^{-1}\overline{E}^2 - \frac{1}{4}N\epsilon^2 = N^{-1}\overline{E}^2 - \frac{1}{4}b\mu^2 c^4$. Neglecting the $\frac{1}{N}$ term in this formula, when $\overline{E}$ is fixed, we obtain the expression for statistical fluctuations typical of gravitating systems: $\overline{(\Delta E)^2}_{bh} = -\frac{1}{4}b\mu^2 c^4$. The well known relation between the mass-energy fluctuations and the behaviour of entropy near the state of thermal equilibrium, $\overline{(\Delta E)^2} = -k\left(\frac{\partial^2 S}{\partial E^2}\right)^{-1}$, leads to the entropy of such a *truncated system*: $\frac{\partial^2 S_{bh}}{\partial E^2} = 4kb^{-1}\mu^{-2}c^{-4}$. Integrating this last equation gives the inverse temperature $\beta_{bh} = 4b^{-1}\mu^{-2}c^{-4}\overline{E}$, where an arbitrary integration constant is fixed to be zero by demanding that a very massive body is also very cold⁴⁻⁶. The entropy is given by the ‘phenomenological’ entropy formula of Bekenstein¹: $S_{bh} = 2kb^{-1}\mu^{-2}\overline{E}^2 + const.$ The model calculation of Hawking² leads to a numerical value of the constant b , $b = \frac{1}{4\pi^2}$. Quite independently of the actual value of the numerical constant b the entropy S_{bh} has a lower bound $S_0 = 2kb^{-1}\mu^{-2}E_0^2 = k\frac{N}{2}$, which depends only on N .

We have seen the emergence of the Bekenstein formula for the black hole entropy from the hypothesis about the microscopic nature of gravitational phenomena³⁻⁴⁻⁸⁻⁹. We shall propose to apply the same argument to the whole Universe. It seems natural to consider $N_U \sim 10^{123}$ gravitational atoms⁴ and apply to them the same physical argument. We obtain the simple estimate⁴ for cosmological constant regarded as the zero-point vacuum energy density $\lambda \sim \frac{\mu^4}{N_U}$. It was shown⁸ that a system of a large number of gravitational atoms described in the framework of the Atomic Theory of Gravitation (the new gravitational mechanics) by a large $N \sim N_U$ matrix of operators³⁻⁴⁻⁸⁻⁹ bears remarkable similarity to the thermal Universe with a nonvanishing positive cosmological constant. It should be noticed that the notion of a statistical ensemble for the observable Universe is justified only after we identify *atoms* whose existence is underlying the totality of phenomena. The Gibbs-Jaynes principle of the maximum of H function⁷ is applicable to a closed system once we postulate the general *Atomic Hypothesis*³⁻³⁻⁴⁻⁵. According to this hypothesis the totality of phenomena should be derived from the properties of space-time-matter atoms, which I prefer to call *gravitational atoms*³⁻⁴⁻⁵⁻⁶. The result described in this short note, which was based on the purely phenomenological considerations, was first derived in the Spring of 1995 and published in⁴. I have applied the

^aTwo examples of gravitating systems are discussed in⁴⁶ simultaneously: a spin-zero quantum black hole and the whole non-rotating Universe. The present upper bound on cosmological constant λ in Planck units, regarded as the vacuum zero-point energy density, was used to estimate the lower bound on the total number $N_U \sim 10^{123}$ of gravitational atoms (and gravitational oscillators) in the Universe.

simple interpolation argument, originally due to Planck, to the problem of mass-energy fluctuations of a gravitating mass^{4 6}. I have proposed that the formalism of **the new gravitational noncommutative mechanics**^{3 4 5 8 9} be applied to gravitating particles (black holes) and to the whole Universe. The point made very clearly in³ was that the microscopic theory of gravitation requires new ideas of the kind Heisenberg has once introduced. The idea of the *Second Heisenberg Algebra* was then introduced^{3 5 8 9}. It is obvious to the present author that this idea is quite useful once we realize that the perihelion precession of the Mercury, and the *Complementarity Principle* for a gravitational mass and a inertial mass aspect of a gravitating particle^{3 5 8 9} allow for the transition to new gravitational mechanics in the same way the nonrelativistic Kepler problem and the spectroscopic data has made it possible for Heisenberg to propose the matrix quantum mechanics³. The idea of using the GRT Kepler problem to make a similar transition to the new gravitational noncommutative mechanics was first described by the present author^{3 5} as early as in the Spring of 1995, and later during the USC Summer 1995 Institute⁵. The idea of what I have called the *Second Heisenberg Algebra* and the *Second Period of Nature* was also described in numerous lectures and talks (CALTECH, February 1996; Tel-Aviv University, March 1996; Ecole Polytechnique, Paris, October 1996). The picture of a gravitating particle (a black hole) which has emerged from my work is not unlike that of a giant nucleus or a baryon.^b The calculation of specific properties of such a complex object like a quantum black hole must be carried out in a sort of large N approximation in the new gravitational noncommutative mechanics proposed in^{3 5 6 8 9}, and it is under way.

Acknowledgments

This research was partially supported by a NSF grant to USC. I wish to thank organizers of the Eight Marcel Grossmann Meeting in Jerusalem for inviting me to this Meeting, which, however, I was unable to attend.

References

1. J. D. Bekenstein, PRD **7**, 2333 (1973).
2. S. W. Hawking, Commun. Math. Phys. **43**, 199 (1975).
3. P. O. Mazur, Acta Phys. Polon. B **26**, 1685 (1995).
4. P. O. Mazur, Acta Phys. Polon. B **27**, 1849 (1996).
5. P. O. Mazur, Lecture at the Summer Institute at USC, 1995.
6. A. Z. Górski and P. O. Mazur, *The Quantum Black Hole Specific Heat is Positive*, hep-th/9704179, submitted to *Phys. Rev. Lett.*.
7. S. P. Heims and E. T. Jaynes, Rev. Mod. Phys. **34**, 143 (1962).
8. P. O. Mazur, *On the Fundamental Idea of the New Gravitational Mechanics*, Spring 1996, unpublished.
9. P. O. Mazur, *Gravitation as a Many Body Problem*, in *Beyond the Standard Model V*, eds. G. Eigen, P. Osland, and B. Stugu, AIP Proceedings 415, pp.

^bThe large N QCD description of baryons and the Wigner (or the modern matrix models) description of energy levels in nuclei share some common traits.

299-305, New York 1997.